

(19) World Intellectual Property
Organization
International Bureau



10 DEC 2004



(43) International Publication Date
31 December 2003 (31.12.2003)

PCT

(10) International Publication Number
WO 2004/000802 A2

(51) International Patent Classification⁷: **C07D**
(21) International Application Number:
PCT/US2003/019779

(US). NI, Jiahong [CN/US]; 505 Castle Drive, Baltimore, MD 21212 (US). LI, Hengguang [CN/US]; 4077 McDowell Lane, Baltimore, MD 21227 (US). SINGH, Suddam [GB/US]; 33 Forest Park, Durham, NH 03824 (US).

(22) International Filing Date: 20 June 2003 (20.06.2003)

(74) Agent: **HULTQUIST, Steven, J.**; Intellectual Property/Technology Law, P.O. Box 14329, Research Triangle Park, NC 27709 (US).

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/390,776 20 June 2002 (20.06.2002) US

(71) Applicant (for all designated States except US): **UNIVERSITY OF MARYLAND BIOTECHNOLOGY INSTITUTE** [US/US]; Suite 200, 701 East Pratt Street, Baltimore, MD 21202-3101 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.


(72) Inventors; and

(75) Inventors/Applicants (for US only): **WANG, Lai-Xi** [CN/US]; 3817 Palmetto Court, Ellicott City, MD 21042

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO,

[Continued on next page]

(54) Title: SCAFFOLDED MALEIMIDE CLUSTERS FOR MULTIVALENT PEPTIDE ASSEMBLY



(57) Abstract: Disclosed are scaffolded maleimide clusters, methods of making said clusters and use of said clusters as templates for multivalent peptide assembly. Multiple maleimide functionalities were introduced onto a scaffold molecule by the reaction of a core-centered polyamines with methoxycarbonyl maleimide or with activated esters of maleimide-containing compounds. The scaffolded maleimides allow rapid, highly chemoselective, and high-yield ligation with thiolcontaining peptides under virtually neutral conditions at room temperature. The disclosed mild and highly efficient ligation method is extremely valuable for synthesizing large and complex multivalent peptides that may not be easily obtained by conventional ligation methods. These novel scaffolded maleimide clusters allow a highly chemoselective ligation with a thiolcontaining peptide under virtually neutral conditions, providing a new and efficient approach for multivalent peptide assembly. The disclosed mild and highly efficient ligation method is extremely valuable for synthesizing large and complex multivalent peptides that may not be easily obtained by conventional ligation methods. A series of multivalent peptides containing the sequence of the 36-mer HIV-1 inhibitor DP178 (T20), the T-helper epitope from tetanus toxoid (830-844), and the minimum epitope sequence of the potent HIV-neutralizing antibody 2F5 were synthesized. Carbohydrates and cholic acid were chosen as the scaffold because of their rigidity

and multi-functionality. Thus, the topology of the multivalent peptides can be controlled by the defined spatial orientation of the maleimide functionalities on the rigid scaffold core. The resulting multivalent gp41 peptides incorporating strands of DP178 on the monosaccharide and the cholic acid templates were found to be able to form three or four α -helix bundles. Moreover, the multivalent peptides containing strands of the long gp41 peptide DP178 were highly immunogenic and were able to raise high titers of peptide-specific antibodies, in the absence of any additional adjuvant. Therefore, these and related multivalent peptides constructed on the maleimide clusters may be used as novel immunogens, potential inhibitors, protein mimics, artificial proteins, and powerful antigens for a broad range of biomedical applications.

WO 2004/000802 A2